esp@cenet document view

Page 1 of 1

PRODUCTION OF RECOMBINANT PROTEIN

JP3244391

1991-10-31

allerat:

YASUEDA HIŞASHI; NAGASE KAZUO

AJINOMOTO KK

C12N15/09; C12N1/21; C12N15/18; C12N15/26; C12P1/00; C12P1/02; C12R1/19; C12P16/09; C12P1/00; C12P1/02; C12R1/19; C12N15/09; C12N1/21; C12N15/16; C12N15/19; C12P21/00; C12P21/02; C12N15/09; (IPC1-7); C12N15/16; C12N15/26; C12P21/00; C12P21/02

m JP19900041666 19900222 Priority number(s): JP19900041666 19900222

Abstract of JP3244391

PURPOSE:To produce a useful protein by culturing Escherichia coli transformed with an expression PURPOSE:To produce a useful protein by culturing Escherichia coll bransformed with an expression plasmid in which a DNA capable of coding the useful protein such as human B cell differentiation factor is integrated in a culture medium containing a soybean hydrolyzate added thereto. CONSTITUTION:An expression plasmid in which a DNA capable of coding a useful protein such as human B cell differentiation factor or human interleukin 2 is integrated is used to transform Escherichia coll. The resultant transformed Escherichia coll is then cultured in a culture medium containing a soybean hydrolyzate added thereto to produce the useful protein, which is subsequently recovered from the culture medium. The amount of the scybean hidrolyzate added to the culture medium is preferably 1-10 wt.% based on the culture medium. The culturing temperature is preferably about 20-45 deg.C, more preferably about 36-39 deg.C in the aforementioned culturing and the pH during the culturing is preferably about 5-9, more preferably about 6-7.

Data supplied from the esp@cenetdatabase - Worldwide